

BROW SURGERY AFTERCARE INSTRUCTIONS

- If you received anesthesia during the procedure, you will be sleepy for the rest of the day and maybe tomorrow. DO NOT drive, operate any machinery, or do anything requiring coordination for the rest of the day. Do not sign any legal documents or make any important decisions in the next 24 hours. After anesthesia, you should breathe deeply each waking hour in order to keep the lungs clear.
- On the day of your surgery and during that night, you should have a responsible adult to be with you or for your protection and safety. You may go to the bathroom with assistance.
- Use medications as instructed. Many patients find Tylenol to be adequate to control postoperative discomfort. DO NOT take aspirin products or NSAIDS (e.g. Ibuprofen, Advil, Aleve, Motrin, Naproxen, Excedrin). If any of these are prescribed to you by another physician, please advise our office.

DIET

- For the first 24 hours after surgery we recommend a light diet. We do not recommend *any alcoholic beverages for 24 hours after your procedure. After that, we do not recommend combining your postoperative pain medication with alcoholic beverages.* After the first 24 hours you may advance to a regular diet as you can tolerate.

YOUR DRESSING

- DO NOT change your dressing or get it wet unless instructed otherwise. You may resume showering below the dressings 2 days after surgery. Please try to keep the dressing dry.
- You will have a head wrap dressing. This dressing will be removed at 24 hours.

THINGS TO HAVE

- Prescription Pain Medication
- Prescription Nausea Medication
- Aquaphor (over the counter)
- Stool Softener (over the counter)

WOUND CARE

- Wound care is extremely important and can help with healing. Please follow the instructions carefully.
- After the dressings are removed if present based on the type of browlift, leave them off. You will still have sutures and staples in place. The sutures/staples will be removed 7-10 days after surgery, or when clinically indicated
You will have sutures and staples in areas of the hair-bearing scalp. Be careful when washing your hair or combing so that the incisions do not separate and the sutures do not get caught. Do not rub forehead or scalp area. This may disrupt the lift.
- Instructions on washing your hair:
 1. Mix Baby shampoo with warm water in a container.
 2. Pour the solution over your hair several times.
 3. Stand under the shower and rinse off.

WHAT IS NORMAL

- You will experience moderate discomfort for 2 to 3 days after surgery.
- Slight oozing of blood from the incision site over the first 2 days.
- Swelling, which is the body's way of healing, can last for several months; the swelling will likely be worse on day 2 or 3 after surgery and then improve significantly over the next several weeks.
- Some black and blueness which is associated with healing; because of gravity, the black and blueness may settle down into your cheeks. All patients are different: sometimes the bruising is gone in less than a week, other times it lasts 2-3 weeks.
- Feeling of dryness (like there is sand in your eye) or excessive tearing; lubricating eye ointment and artificial tears at least 4 to 6 times a day will help this.
- Some blurry vision from swelling and ointment.
- Take comfort in the fact that the "tight" feeling, numbness, bumps and irregularities, are normal after surgery and slowly improve over months.
- The body heals a surgical scar in slow stages. Gradually improving surgical scars and contours can be expected **for up to 1 year after surgery**.
- Temporary hair loss along the incision lines.
- As the bruising and swelling start to decrease and you start to pick up some more of your normal routine, you will feel more like your normal self.

WHAT TO AVOID

- Driving or operating heavy equipment after sedation or while taking prescription pain medication.
- Heavy lifting or strenuous activities for 7 days.
- Makeup until the sutures are removed or the incision lines are well healed.
- Sun exposure; if you must go into the sun, wear a large brimmed hat and sunscreen.
- Sleeping flat. Place 2-3 pillows behind head at night as it will help with swelling.
- Smoking/tobacco as these slow the healing process.

AFTERCARE ACTIVITY

- You may resume showering and shampooing 2 days after your surgery date. Try to keep the dressing dry.
- The **ONLY** exercise for the first 3 weeks is walking: Outside walking is ok after 2wks.
- You may return to the gym after 4wks, doing only one-half of your usual regimen and take 3 weeks to return to your regular regimen. Weight, machines, running, swimming, golf, tennis, yoga, and Pilates after 3 weeks
- Please refrain from sexual activity for 3-4 weeks.
- For the first 3 weeks lift no more than 10 pounds.
- You may return to sedentary work 2 weeks after your surgery.
- Please refrain from driving a car for at least 3-5 days after your surgery date. After 3-5 days you may resume driving if you feel comfortable. Please ensure you are off all narcotic pain medications before resuming driving.

Call Dr. Patel on his cell phone (253)733-7048 if you experience the following:

- A sudden, severe decrease or dimness in your vision.
- Severe, unrelenting, or stabbing pains, particularly if it is associated with vomiting or fever.
- Extreme swelling around the surgical area that is very hot, firm, or tense.
- If you have severe or increased pain not relieved by medication.
- If you have any side effects to medication; such as, rash, nausea, headache, vomiting.
- If you have an oral temperature over 101 degrees.

- If you have any yellowish or greenish drainage or notice a foul odor.
- If you have (excessive) bleeding or saturation of the dressing. A small amount of bleeding may appear on the dressing or pad.
- If you have an increase in pain after the first 48 hours.
- If you experience persistent nausea or vomiting.

If you experience a medical emergency and are unable to reach the doctor on call, go to the nearest emergency room.