

LIPOSUCTION WITH RENUVION AFTERCARE INSTRUCTIONS

- If you received anesthesia during the procedure, you will be sleepy for the rest of the day and maybe tomorrow. **DO NOT** drive, operate any machinery, or do anything requiring coordination for the rest of the day. Do not sign any legal documents or make any important decisions in the next 24 hours. If you are a smoker, please do not smoke for 2-3wks. It will interfere with wound healing. After anesthesia, you should breathe deeply each waking hour in order to keep the lungs clear.
- On the day of your surgery and during that night, you should have a responsible adult to be with you for your safety. You may need assistance to go to the bathroom.
- Use medications as instructed. Patients may find Tylenol to be adequate to control postoperative discomfort if they prefer not to take any narcotic medications. **DO NOT** take aspirin products or nsaid (e.g. Ibuprofen, Advil, Aleve, Motrin, Naproxen, Excedrin). If any of these are prescribed to you by another physician, please advise our office.

DIET

- For the first 24 hours after surgery we recommend a light diet. We do not recommend *any alcoholic beverages for 24 hours after your procedure. After that, we do not recommend combining your postoperative pain medication with alcoholic beverages.* After the first 24 hours you may advance to a regular diet as you can tolerate.
- Drink plenty of fluids to stay hydrated.

THINGS TO HAVE

- Prescription Pain Medications
- Prescription Nausea Medications
- Stool Softeners (over the counter)

YOUR DRESSING

- Please remove dressings and abdominal binder (if present) after 24 hours if **NO** other surgeries were done at the same time. If you have clear tape on any part of your body, please leave in place until your first postop appointment, unless otherwise advised by Dr. Patel. You may resume showering daily at this point.
- After 24hours you may transition to wearing your compression garments for 4 weeks. You may wash the garments as needed. Additional garments can be purchased at the Cascade Cosmetic centers.

WOUND CARE

- Wound care is extremely important and can help with healing. Please follow the instructions carefully.
- Do not apply ice or heat to the surgical area(s).
- Keep incision clean and dry.
- Do not apply any lotions or ointments without discussing it with Dr. Patel.
- Shower daily after 24hrs.
- Drainage from your incision sites may occur following your procedure. It is recommended to change absorbent pads every 4-6 hours as needed, secondary to drainage which is normal.

WHAT IS NORMAL

- You will experience moderate discomfort for 2 to 3 days after surgery.
- You may experience mild to moderate swelling, bruising, or soreness for 1-2 weeks following your procedure.
- Slight oozing of blood and moderate “clear to fruit punch” colored drainage from the incision sites over the first 2-3days is common.
- You may feel excess gas in your tissues (subcutaneous air), this is normal and will subside in 2-3 days.
- Take comfort in the fact that the “tight” feeling, numbness, bumps and irregularities, are normal after surgery and slowly improve over months.
- The body heals a surgical scar in slow stages. Gradually improving surgical scars and contours can be expected **for up to 1 year after surgery.**

WHAT TO AVOID

- Driving or operating heavy equipment after sedation or while taking prescription pain medication.
- Heavy lifting (greater than 10 pounds) or strenuous activities for 2-3 weeks.
- Sleeping flat. Place 2-3 pillows behind head at night as it will help with discomfort.
- Smoking/tobacco as these slow the healing process.

AFTERCARE ACTIVITY

- You may resume showering 1 day after your surgery date.
- You may return to regular activities after 1 week.
- You may return to the gym after 1 week
- Please refrain from sexual activity for 1 week.
- You may return to sedentary work 2 days after your surgery, if you are no longer taking narcotic pain medications.
- Please refrain from driving a car for at least 2 days after your surgery date. Please ensure you are off all narcotic pain medications before resuming driving.

Call Dr. Patel on his cell phone (253)733-7048 if you experience the following:

- Extreme swelling around the surgical area that is very hot, firm, or tense.

- If you have severe or increased pain not relieved by medication.
- If you have any side effects to medication; such as, rash, nausea, headache, vomiting.
- If you have an oral temperature over 101.5 degrees.
- If you have any purulent drainage or notice a foul odor.
- If you have (excessive) bleeding or saturation of the dressing. A small amount of bleeding may appear on the dressing or pad.
- If you have an increase in pain or redness after the first 48 hours.
- If you experience persistent nausea or vomiting.

If you experience a medical emergency and are unable to reach the doctor on call, go to the nearest emergency room.